

CAN DHIVY HELP ME DIVIDE TO CONQUER THE DAY?



Start the conversation about DHIVY, a carbidopa/levodopa (CD/LD) tablet that was made to be broken

Each day with Parkinson's disease can present new challenges. **It's important to ask yourself if your current treatment is meeting your needs day to day, and to be open with your doctor about how you're feeling.**

	STRONGLY AGREE	SOMEWHAT AGREE	SOMEWHAT DISAGREE	STRONGLY DISAGREE
Does your current daily dose of CD/LD help you manage your motor symptoms?				
How often do you experience involuntary movements (also known as dyskinesia), such as twitches, jerks, twisting, or writhing?	DAILY	WEEKLY	MONTHLY	NEVER
How often do you experience your medication "wearing off" (also known as OFF time) or periods when symptoms are not under control?				
How often do you feel that your symptoms are under control and you're maintaining a strong ON time?				
How often do any of the symptoms mentioned above change, impact, or prevent you from doing the activities you want to do?				
How often do you "split" or purposefully break CD/LD tablets to adjust your dose?				

After answering these questions, print this document and bring it to your doctor to ask if DHIVY is right for you.

INDICATION

DHIVY is a prescription medication that contains a combination of carbidopa and levodopa for the treatment of Parkinson's disease, Parkinson's disease caused by infection or inflammation of the brain, or Parkinson's disease-like symptoms that may result from carbon monoxide or manganese poisoning.

IMPORTANT SAFETY INFORMATION

DHIVY is contraindicated in patients

- Currently taking a nonselective monoamine oxidase (MAO) inhibitor (e.g., phenelzine, linezolid, and tranylcypromine) or have recently (within 2 weeks) taken a nonselective MAO inhibitor. Hypertension can occur if these drugs are used concurrently.
- With known hypersensitivity to any component of DHIVY.

Please see Important Safety Information continued on the next page and full Prescribing Information.

Dhivy[®]
(carbidopa/levodopa)
Segments 6.25 mg/25 mg | Tablets 25 mg/100 mg



WHY DHIVY?

Finding the right daily dose of CD/LD can be a challenge.

Your doctor may recommend dividing your daily dose of CD/LD.

However, splitting tablets that aren't designed to be broken could lead to inaccurate doses.

Dhivy[®]
(carbidopa/levodopa)
Segments 6.25 mg/25 mg | Tablets 25 mg/100 mg

DHIVY is a CD/LD tablet designed to be divided.



Image does *not* represent the actual size of the tablet.

IMPORTANT SAFETY INFORMATION (continued)

Taking DHIVY may result in falling asleep while engaged in normal activities, even without warning and as late as 1 year or later after starting DHIVY. This may affect your ability to drive or operate machinery, resulting in accidents. Do not do anything that requires alertness until you know how DHIVY affects you. Tell your healthcare provider about any sleep medicines you take or if you have a sleep disorder, and discuss any episodes of drowsiness or sleepiness you experience while taking DHIVY.

Talk to your healthcare provider before you change the dose or stop taking DHIVY, as this may result in serious side effects. Call your healthcare provider immediately if you develop withdrawal symptoms such as fever, confusion, or severe muscle stiffness.

Tell your healthcare provider if you have any heart conditions, especially if you have had a heart attack or irregular heartbeats. Your heart function should be monitored during the time period your DHIVY dose is first adjusted.

Treatment with DHIVY may cause hallucinations or abnormal thoughts and behaviors. Be sure to tell your healthcare provider if you experience excessive suspicion; seeing, hearing, or feeling things that are not real; confusion; agitation; aggressive behavior; and disorganized thinking.

Treatment with DHIVY may also cause intense urges to gamble, increased sexual urges, intense urges to spend money, binge eating, and other intense urges, as well as the inability to control those urges. Uncontrolled sudden movements of the face, arms, legs, or trunk may appear or get worse during treatment with DHIVY. This may mean your dosage of DHIVY or other Parkinson's medications needs to be adjusted.

Tell your healthcare provider if you have ever had a peptic ulcer or glaucoma, as treatment with DHIVY may worsen these conditions. DHIVY also may cause abnormal blood or urine test results.

Some patients taking DHIVY may experience depression or suicidal thoughts. Tell your healthcare provider if you have thoughts of suicide.

The most common side effects that may appear with DHIVY include uncontrolled sudden movements and nausea.

Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using DHIVY with certain other medicines, including MAO inhibitors, isoniazid, iron supplements, medicines for high blood pressure or those that increase or decrease dopamine levels, or metoclopramide, may cause serious side effects.

Notify your healthcare provider if you become pregnant or intend to become pregnant during DHIVY therapy or if you intend to breastfeed or are breastfeeding an infant.

Please see [full Prescribing Information](#).